Neuroscience Letters, 167 (1994) 198–200
Elsevier Science Ireland Ltd.

AUTHOR INDEX

Abraham, W.C., see Christie, B.R., 41
Arai, A., see del Cerro, S., 149
Araki, W., Kunishita, T., Takahashi, K., Ikeda, S. and Tabira, T., Demonstration of amyloid β-protein secretion in a mouse neuronal cell line, 125
Arendt, J., see Deacon, S., 191
Arita, J., see Izaki, Y., 171
Arregui, A., Hollingsworth, Z., Penney, J.B. and Young, A.B., Autoradiographic evidence for increased dopamine uptake sites in striatum of hypoxic mice, 195
Arvidsson, J., see Johansson, K., 187
Averill, S., see Ching, Y.P., 67
Avidor, R., see Gozes, I., 24
Avoli, M., see Louvel, J., 97
Axelrad, H., see Berthié, B., 161

Bahr, B.A., see del Cerro, S., 149
Baptista, T., Weiss, S.R.B., Zocchi, A., Sitcoske, M. and Post, R., Electrical kindling is associated with increases in amygdala acetylcholine levels: an in vivo microdialysis study, 133
Barbagli, B., see Buda, M., 33
Barnard, E.A., see Matsumoto, M., 183
Basham, D.A., see Stone, E.A., 11
Bäurle, J. and Grüsser-Cornehls, U., Calbindin D-28k in the lateral vestibular nucleus of mutant mice as a tool to reveal Purkinje cell plasticity, 85
Behrmann, G., see Pritz-Hohmeier, S., 59
Berthié, B. and Axelrad, H., Granular layer collaterals of the unipolar brush cell axon display rosette-like excrescences. A Golgi study in the rat cerebellar cortex, 161
Bette, M., see Schäfer, M.K.-H., 137
Bileviciute, I., Lundeberg, T., Ekblom, A. and Theodorsson, E., Substance P-, neurokinin A-, calcitonin gene-related peptide- and neuropeptide Y-like immunoreactivity (-LI) in rat knee joint synovial fluid during acute monoarthritis is not correlated with concentrations of neuropeptide-LI in cerebrospinal fluid and plasma, 145
Bing, G., see Stone, E.A., 11
Blizard, D., see Buda, M., 33
Borbély, A.A., see Franken, P., 89
Buda, M., Lachuer, J., Devauges, V., Barbagli, B., Blizard, D. and Sara, S.J., Central noradrenergic reactivity to stress in Maudsley rat strains, 33

Callera, J.C., Saad, W.A., De Arruda Camargo, L.A., Renzi, A., De Luca, Jr., L.A. and Menani, J.V., Role of the adrenergic pathways of the lateral hypothalamus on water intake and pressor response induced by the cholinergic activation of the medial septal area in rats, 153
Chang, A.C., see Chang, N.-C., 105
Chang, C.-L., see Cheng, J.-T., 117
Chang, N.-C., Jenkins, N.A., Gilbert, D.J., Copeland, N.G., Chang, Y.-H., Chen, W.-M. and Chang, A.C., Assignment of two α2 adrenoceptor subtype genes to murine chromosomes, 105
Chang, Y.-H., see Chang, N.-C., 105
Chen, W.-M., see Chang, N.-C., 105
Cheng, J.-T., Chang, C.-L. and Tsai, C.-L., Inhibitory effect of neuropeptide Y (NPY) on the in vitro activity of tyrosine hydroxylase, 117
Ching, Y.P., Averill, S., Wilkin, G.P., Wotherspoon, G. and Priestley, J.V., Serotonergic terminals express a growth associated protein (GAP-43) in the adult rat spinal cord, 67
Choi, D.W., see Weiss, J.H., 93
Christie, B.R. and Abraham, W.C., L-type voltage-sensitive calcium channel antagonists block heterosynaptic long-term depression in the dentate gyrus of anaesthetized rats, 41
Cline, E.J., see Orona, R.A., 121
Copeland, N.G., see Chang, N.-C., 105

Davidson, B.L., see Roessler, B.J., 5
De Arruda Camargo, L.A., see Callera, J.C., 153
de Boer, P., see Westerink, B.H.C., 109
De Luca, Jr., L.A., see Callera, J.C., 153
De Vries, J.B., see Westerink, B.H.C., 109
Deacon, S. and Arendt, J., Posture influences melatonin concentrations in plasma and saliva in humans, 191
del Cerro, S., Arai, A., Kessler, M., Bahr, B.A., Vanderklish, P., Rivera, S. and Lynch, G., Stimulation of NMDA receptors activates calpain in cultured hippocampal slices, 149
Delivoria-Papadopoulos, M., see Hoffman, D.J., 156
Devauges, V., see Buda, M., 33
Dijk, D.-J., see Franken, P., 89
Dimlich, R.V.W., see Medrano, S., 14
Dobson, S.P., Quinn, J.P., Morrow, J.A. and Mulderry, P.K., The rat vasoactive intestinal polypeptide cyclic AMP response element regulates gene transcriptional responses differently in neonatal and adult rat sensory neurons, 19
Dussaillant, M., see Gozes, I., 24
Dymshitz, J. and Vasko, M.R., Endothelin-1 enhances capsaicin-induced peptide release and cGMP accumulation in cultures of rat sensory neurons, 128

Ekblom, A., see Bileviciute, I., 145

Farkas, T., see Toldi, J., 1
Ferres-Torres, R., see González-Hernández, T., 77
Fonnum, F., see Hassel, B., 29
Franken, P., Dijk, D.-J., Tobler, I. and Borbély, A.A., High-frequency components of the rat electrocorticogram are modulated by the vigilance states, 89

Galligan, J.J., see Yunker, A.M., 51
Galvan, I., see Pranzatelli, M.R., 166
Giladi, E., see Gozes, I., 24
Gilbert, D.J., see Chang, N.-C., 105
Golabek, A.A., see Kida, E., 73
González-González, B., see González-Hernández, T., 77
González-Hernández, T., Mantolán-Sarmiento, B., González-González, B., Ferres-Torres, R. and Meyer, G., NADPH-d (dihydronicotinamide adenine dinucleotide phosphate diaphorase) activity in geniculo-tectal neurons of the rat, 77
Gozes, I., Avidor, R., Giladi, E., Shani, Y., McEwen, B.S., Dussaillant, M. and Rostene, W., Adrenalectomy decreases vasoactive intestinal peptide mRNA levels in the rat suprachiasmatic nucleus, 24
Grolleau, F. and Lapied, B., Transient Na^+-activated K^+ current in beating pacemaker-isolated adult insect neurosecretory cells (dum neurones), 46

Neuroscience Letters, 167 (1994) 198–200
Elsevier Science Ireland Ltd.

AUTHOR INDEX

Abraham, W.C., see Christie, B.R., 41
Arai, A., see del Cerro, S., 149
Araki, W., Kunishita, T., Takahashi, K., Ikeda, S. and Tabira, T., Demonstration of amyloid β-protein secretion in a mouse neuronal cell line, 125
Arendt, J., see Deacon, S., 191
Arita, J., see Izaki, Y., 171
Arregui, A., Hollingsworth, Z., Penney, J.B. and Young, A.B., Autoradiographic evidence for increased dopamine uptake sites in striatum of hypoxic mice, 195
Arvidsson, J., see Johansson, K., 187
Averill, S., see Ching, Y.P., 67
Avidor, R., see Gozes, I., 24
Avoli, M., see Louvel, J., 97
Axelrad, H., see Berthié, B., 161

Bahr, B.A., see del Cerro, S., 149
Baptista, T., Weiss, S.R.B., Zocchi, A., Sitcoske, M. and Post, R., Electrical kindling is associated with increases in amygdala acetylcholine levels: an in vivo microdialysis study, 133
Barbagli, B., see Buda, M., 33
Barnard, E.A., see Matsumoto, M., 183
Basham, D.A., see Stone, E.A., 11
Bäurle, J. and Grüsser-Cornehls, U., Calbindin D-28k in the lateral vestibular nucleus of mutant mice as a tool to reveal Purkinje cell plasticity, 85
Behrmann, G., see Pritz-Hohmeier, S., 59
Berthié, B. and Axelrad, H., Granular layer collaterals of the unipolar brush cell axon display rosette-like excrescences. A Golgi study in the rat cerebellar cortex, 161
Bette, M., see Schäfer, M.K.-H., 137
Bileviciute, I., Lundeberg, T., Ekblom, A. and Theodorsson, E., Substance P-, neurokinin A-, calcitonin gene-related peptide- and neuropeptide Y-like immunoreactivity (-LI) in rat knee joint synovial fluid during acute monoarthritis is not correlated with concentrations of neuropeptide-LI in cerebrospinal fluid and plasma, 145
Bing, G., see Stone, E.A., 11
Blizard, D., see Buda, M., 33
Borbély, A.A., see Franken, P., 89
Buda, M., Lachuer, J., Devauges, V., Barbagli, B., Blizard, D. and Sara, S.J., Central noradrenergic reactivity to stress in Maudsley rat strains, 33

Callera, J.C., Saad, W.A., De Arruda Camargo, L.A., Renzi, A., De Luca, Jr., L.A. and Menani, J.V., Role of the adrenergic pathways of the lateral hypothalamus on water intake and pressor response induced by the cholinergic activation of the medial septal area in rats, 153
Chang, A.C., see Chang, N.-C., 105
Chang, C.-L., see Cheng, J.-T., 117
Chang, N.-C., Jenkins, N.A., Gilbert, D.J., Copeland, N.G., Chang, Y.-H., Chen, W.-M. and Chang, A.C., Assignment of two α2 adrenoceptor subtype genes to murine chromosomes, 105
Chang, Y.-H., see Chang, N.-C., 105
Chen, W.-M., see Chang, N.-C., 105
Cheng, J.-T., Chang, C.-L. and Tsai, C.-L., Inhibitory effect of neuropeptide Y (NPY) on the in vitro activity of tyrosine hydroxylase, 117
Ching, Y.P., Averill, S., Wilkin, G.P., Wotherspoon, G. and Priestley, J.V., Serotonergic terminals express a growth associated protein (GAP-43) in the adult rat spinal cord, 67
Choi, D.W., see Weiss, J.H., 93
Christie, B.R. and Abraham, W.C., L-type voltage-sensitive calcium channel antagonists block heterosynaptic long-term depression in the dentate gyrus of anaesthetized rats, 41
Cline, E.J., see Orona, R.A., 121
Copeland, N.G., see Chang, N.-C., 105

Davidson, B.L., see Roessler, B.J., 5
De Arruda Camargo, L.A., see Callera, J.C., 153
de Boer, P., see Westerink, B.H.C., 109
De Luca, Jr., L.A., see Callera, J.C., 153
De Vries, J.B., see Westerink, B.H.C., 109
Deacon, S. and Arendt, J., Posture influences melatonin concentrations in plasma and saliva in humans, 191
del Cerro, S., Arai, A., Kessler, M., Bahr, B.A., Vanderklish, P., Rivera, S. and Lynch, G., Stimulation of NMDA receptors activates calpain in cultured hippocampal slices, 149
Delivoria-Papadopoulos, M., see Hoffman, D.J., 156
Devauges, V., see Buda, M., 33
Dijk, D.-J., see Franken, P., 89
Dimlich, R.V.W., see Medrano, S., 14
Dobson, S.P., Quinn, J.P., Morrow, J.A. and Mulderry, P.K., The rat vasoactive intestinal polypeptide cyclic AMP response element regulates gene transcriptional responses differently in neonatal and adult rat sensory neurons, 19
Dussaillant, M., see Gozes, I., 24
Dymshitz, J. and Vasko, M.R., Endothelin-1 enhances capsaicin-induced peptide release and cGMP accumulation in cultures of rat sensory neurons, 128

Ekblom, A., see Bileviciute, I., 145

Farkas, T., see Toldi, J., 1
Ferres-Torres, R., see González-Hernández, T., 77
Fonnum, F., see Hassel, B., 29
Franken, P., Dijk, D.-J., Tobler, I. and Borbély, A.A., High-frequency components of the rat electrocorticogram are modulated by the vigilance states, 89

Galligan, J.J., see Yunker, A.M., 51
Galvan, I., see Pranzatelli, M.R., 166
Giladi, E., see Gozes, I., 24
Gilbert, D.J., see Chang, N.-C., 105
Golabek, A.A., see Kida, E., 73
González-González, B., see González-Hernández, T., 77
González-Hernández, T., Mantolán-Sarmiento, B., González-González, B., Ferres-Torres, R. and Meyer, G., NADPH-d (dihydronicotinamide adenine dinucleotide phosphate diaphorase) activity in geniculo-tectal neurons of the rat, 77
Gozes, I., Avidor, R., Giladi, E., Shani, Y., McEwen, B.S., Dussaillant, M. and Rostene, W., Adrenalectomy decreases vasoactive intestinal peptide mRNA levels in the rat suprachiasmatic nucleus, 24
Grolleau, F. and Lapied, B., Transient Na^+-activated K^+ current in beating pacemaker-isolated adult insect neurosecretory cells (dum neurones), 46

Gruenstein, E., see Medrano, S., 14
Grüsser-Cornehls, U., see Bäurle, J., 85

Ham, J. and Scanlon, M., The glutamate inhibition of carbachol stimulated inositol phosphate production in rat cortical cells is mediated through an ionotropic NMDA receptor, 63
Härtig, W., see Pritz-Hohmeier, S., 59
Hashimoto, M., see Izaki, Y., 171
Hassel, B., Iversen, E.G. and Fonnum, F., Neurotoxicity of albumin in vivo, 29
Hay, M. and Kunze, D.L., An intermediate conductance calcium-activated potassium channel in rat visceral sensory afferent neurons, 179
Hibino, H., see Izaki, Y., 171
Hoffman, D.J., McGowan, J.E., Marro, P.J., Mishra, O.P. and Delivoria-Papadopoulos, M., Hypoxia-induced modification of the *N*-methyl-D-aspartate receptor in the brain of the newborn piglet, 156
Hollingsworth, Z., see Arregui, A., 195
Holsboer, F., see Skutella, T., 55

Ikawa, K., Watanabe, A., Motohashi, N. and Kaneno, S., The effect of repeated administration of methamphetamine on dopamine uptake sites in rat striatum, 37
Ikeda, S., see Araki, W., 125
Iriki, M., see Izaki, Y., 171
Iversen, E.G., see Hassel, B., 29
Izaki, Y., Hashimoto, M., Arita, J., Iriki, M. and Hibino, H., Intraperitoneal injection of 1-oleoyl-2-docosahexaenoyl phosphatidylcholine enhances discriminatory shock avoidance learning in rats, 171

Jenkins, N.A., see Chang, N.-C., 105
Jirikowski, G.F., see Skutella, T., 55
Johansson, K. and Arvidsson, J., Central plasticity in rat trigeminal primary sensory neurons innervating vibrissae after neonatal peripheral nerve injury, 187

Kaneno, S., see Ikawa, K., 37
Kawai, Y., see Umemoto, S., 101
Kessler, M., see del Cerro, S., 149
Kida, E., Golabek, A.A., Wisniewski, T. and Wisniewski, K.E., Regional differences in apolipoprotein E immunoreactivity in diffuse plaques in Alzheimer's disease brain, 73
Kimura, I., see Nojima, H., 113
Kimura, M., see Nojima, H., 113
Kunishita, T., see Araki, W., 125
Kunze, D.L., see Hay, M., 179
Kurcewicz, I., see Louvel, J., 97

Lachuer, J., see Buda, M., 33
Lapied, B., see Grolleau, F., 46
Louvel, J., Avoli, M., Kurcewicz, I. and Pumain, R., Extracellular free potassium during synchronous activity induced by 4-aminopyridine in the juvenile rat hippocampus, 97
Lundeberg, T., see Bileviciute, I., 145
Lynch, G., see del Cerro, S., 149

Manavalan, J.S., see Stone, E.A., 11
Mantolán-Sarmiento, B., see González-Hernández, T., 77
Marro, P.J., see Hoffman, D.J., 156
Matsumoto, M. and Barnard, E.A., Effects of an activator and an inhibitor of protein kinase C on serotonin receptors induced by rat brain mRNA in *Xenopus* oocytes, 183
Mayfield, R.D., see Orona, R.A., 121
McEwen, B.S., see Gozes, I., 24
McGowan, J.E., see Hoffman, D.J., 156
Medrano, S., Gruenstein, E. and Dimlich, R.V.W., Substance P receptors on human astrocytoma cells are linked to glycogen breakdown, 14
Menani, J.V., see Callera, J.C., 153
Meyer, G., see González-Hernández, T., 77
Minamoto, Y., Tanaka, T., Shibata, S. and Watanabe, S., Neuroprotective effect of cholecystokinin$_B$ receptor antagonist on ischemia-induced decrease in CA1 presynaptic fiber spikes in rat hippocampal slices, 81
Mishra, O.P., see Hoffman, D.J., 156
Morrisett, R.A., Potentiation of *N*-methyl-D-aspartate receptor-dependent afterdischarges in rat dentate gyrus following in vitro ethanol withdrawal, 175
Morrow, J.A., see Dobson, S.P., 19
Motohashi, N., see Ikawa, K., 37
Mulderry, P.K., see Dobson, S.P., 19

Noguchi, K., see Umemoto, S., 101
Nojima, H., Kimura, I. and Kimura, M., The evidence of accelerative interaction between cAMP-dependent protein kinase and external calcium for the desensitization of nicotinic acetylcholine receptor channel in mouse skeletal muscle cells, 113

Orona, R.A., Mayfield, R.D., Cline, E.J. and Zahniser, N.R., Repeated intravenous cocaine administration to rats produces behavioral sensitization without changing brain cocaine levels, 121

Penney, J.B., see Arregui, A., 195
Post, R., see Baptista, T., 133
Pranzatelli, M.R. and Galvan, I., Ontogeny of [^{125}I]iodocyanopindolol-labelled 5-hydroxytryptamine$_{1B}$-binding sites in the rat CNS, 166
Priestley, J.V., see Ching, Y.P., 67
Pritz-Hohmeier, S., Härtig, W., Behrmann, G. and Reichenbach, A., Immunocytochemical demonstration of astrocytes and microglia in the whale brain, 59
Probst, J.C., see Skutella, T., 55
Pumain, R., see Louvel, J., 97

Quinn, J.P., see Dobson, S.P., 19

Reichenbach, A., see Pritz-Hohmeier, S., 59
Renzi, A., see Callera, J.C., 153
Reymann, K.G., see Riedel, G., 141
Riedel, G., Wetzel, W. and Reymann, K.G., (*R,S*)-α-Methyl-4-carboxyphenylglycine (MCPG) blocks spatial learning in rats and long-term potentiation in the dentate gyrus in vivo, 141
Rivera, S., see del Cerro, S., 149
Roessler, B.J. and Davidson, B.L., Direct plasmid mediated transfection of adult murine brain cells in vivo using cationic liposomes, 5
Romeo, H., see Schäfer, M.K.-H., 137
Rostene, W., see Gozes, I., 24

Saad, W.A., see Callera, J.C., 153
Santiago, M., see Westerink, B.H.C., 109
Sara, S.J., see Buda, M., 33
Scanlon, M., see Ham, J., 63
Schäfer, M.K.-H., Bette, M., Romeo, H., Schwaeble, W. and Weihe,

E., Localization of κ-opioid receptor mRNA in neuronal subpopulations of rat sensory ganglia and spinal cord, 137
Schwaeble, W., see Schäfer, M.K.-H., 137
Senba, E., see Umemoto, S., 101
Shani, Y., see Gozes, I., 24
Shibata, S., see Minamoto, Y., 81
Sitcoske, M., see Baptista, T., 133
Skutella, T., Probst, J.C., Jirikowski, G.F., Holsboer, F. and Spanagel, R., Ventral tegmental area (VTA) injections of tyrosine hydroxylase phosphorothioate antisense oligonucleotide suppress operant behavior in rats, 55
Spanagel, R., see Skutella, T., 55
Stone, E.A., Manavalan, J.S., Basham, D.A. and Bing, G., Effect of yohimbine on nerve growth factor mRNA and protein levels in rat hippocampus, 11

Tabira, T., see Araki, W., 125
Takahashi, K., see Araki, W., 125
Tanaka, T., see Minamoto, Y., 81
Theodorsson, E., see Bileviciute, I., 145
Tobler, I., see Franken, P., 89
Toldi, J., Farkas, T. and Völgyi, B., Neonatal enucleation induces cross-modal changes in the barrel cortex of rat. A behavioural and electrophysiological study, 1
Tsai, C.-L., see Cheng, J.-T., 117
Turetsky, D., see Weiss, J.H., 93

Umemoto, S., Noguchi, K., Kawai, Y. and Senba, E., Repeated stress reduces the subsequent stress-induced expression of Fos in rat brain, 101

Vanderklish, P., see del Cerro, S., 149
Vasko, M.R., see Dymshitz, J., 128
Völgyi, B., see Toldi, J., 1

Watanabe, A., see Ikawa, K., 37
Watanabe, S., see Minamoto, Y., 81
Weihe, E., see Schäfer, M.K.-H., 137
Weiss, J.H., Turetsky, D., Wilke, G. and Choi, D.W., AMPA/kainate receptor-mediated damage to NADPH-diaphorase-containing neurons is Ca^{2+} dependent, 93
Weiss, S.R.B., see Baptista, T., 133
Westerink, B.H.C., de Boer, P., Santiago, M. and De Vries, J.B., Do nerve terminals and cell bodies of nigrostriatal dopaminergic neurons of the rat contain similar receptors?, 109
Wetzel, W., see Riedel, G., 141
Wilke, G., see Weiss, J.H., 93
Wilkin, G.P., see Ching, Y.P., 67
Wisniewski, K.E., see Kida, E., 73
Wisniewski, T., see Kida, E., 73
Wotherspoon, G., see Ching, Y.P., 67

Young, A.B., see Arregui, A., 195
Yunker, A.M. and Galligan, J.J., Extrinsic denervation increases NADPH diaphorase staining in myenteric nerves of guinea pig ileum, 51

Zahniser, N.R., see Orona, R.A., 121
Zocchi, A., see Baptista, T., 133